

WATER SOLUBLE POWDER

MOXI 50% Powder

DESCRIPTION AND COMPOSITION:

Moxi 50% Powder is a white or yellowish white crystalline powder. Each gram contains 500mg amoxicillin (as trihydrate) as its active ingredient.

TARGET SPECIES:

Chickens, ducks, turkeys and pigs.

INDICATIONS:

Chickens: For the treatment of Pasteurellosis and Colibacillosis.

Turkeys: For the treatment of Pasteurellosis.

Ducks: For the treatment of infections caused by *Streptococcus bovis*, *Pasteurella antipestifer* and *Escherichia coli*.

Pigs: For the treatment of Salmonellosis and Pasteurellosis.

ROUTE OF ADMINISTRATION:

By oral administration via drinking water.

DOSAGE:

Chickens: Administer in drinking water to give 15mg amoxicillin trihydrate per kg body weight. Moxi 50% Powder medicated drinking water should be provided on alternate days. The total period of treatment should be for 3 days (2 days of medication) or in severe cases for 5 days (3 days of medication). This is equivalent to 0.25g to 0.5g of Moxi 50% Powder per litre of drinking water.

Ducks: Administer in drinking water to give 20mg amoxicillin trihydrate per kg body weight. Medication should be provided on alternate days for 3 days (2 days of medication).

Turkeys: Administer in drinking water to give 15mg to 20mg amoxicillin trihydrate per kg body weight. Medication to be provided on alternate days for 5 days (medication on days 1, 3 and 5).

Pigs: Administer in drinking water to give 20mg per kg body weight daily. The dose should be divided and administered at approximately 12 hourly intervals for up to 5 days.

WITHDRAWAL PERIODS:

Meat:

Chickens: 1 day

Ducks: 9 days

Turkeys: 5 days

Pigs: 2 days


Not authorised for use in laying birds producing eggs for human consumption.

SHELF LIFE:

Three (3) years from the date of manufacture.

PACK SIZE:

100g.



Manufacturer: RANGE PHARMA SDN. BHD.
No.1, Jalan TSB 11, Taman Industri Sg. Buloh,
47000 Sg. Buloh, Selangor, Malaysia.
Tel: (603) 6156 8708 Fax: (603) 6156 8707
Email: info@rangepharma.com
Website: www.rangepharma.com

MOXI 50%

粉末

说明和成分:

Moxi 50%粉末是一种白色或淡黄色的结晶粉末。每克含有500毫克amoxicillin（三水合物）为其活性成分。

目标:

鸡, 鸭, 火鸡和猪

适应症:

鸡: 用于治疗Pasteurellosis和Colibacillosis。

火鸡: 用于治疗Pasteurellosis。

鸭子: 用于治疗由Streptococcus bovis, Pasteurella antipestifer和Escherichia coli引起的感染。

猪: 用于治疗Salmonellosis和Pasteurellosis。

用法:

通过饮用水口服

剂量:

鸡: 通过饮用水给药, 每公斤体重摄入15毫克 amoxicillin trihydrate。Moxi 50%粉末含药饮用水应在隔天提供。总治疗期应为3天(用药2天), 或在严重情况下应为5天(用药3天)。这相当于每升饮用水含0.25克至0.5g克的Moxi 50%粉末。

鸭子: 通过饮用水, 每公斤体重摄入20毫克 amoxicillin trihydrate。应在隔天提供药物, 持续3天(服药2天)。

火鸡: 通过饮用水, 每公斤体重摄入15毫克至20毫克amoxicillin trihydrate。应在隔天提供药物, 持续5天(第1、3和5天服药)。

猪: 通过饮用水, 每天以每公斤体重摄入20毫克。剂量应当分配好, 并大约每12小时间隔服用一次, 持续5天。

撤回日期:

肉:

鸡: 1日

鸭子: 9日

火鸡: 5日

猪: 2日


未获准用于产蛋供人类食用的禽类

保质期:

自生产日起三(3)年

包装:

100克



Manufacturer: RANGE PHARMA SDN. BHD.
No.1, Jalan TSB 11, Taman Industri Sg. Buloh,
47000 Sg. Buloh, Selangor, Malaysia.
Tel: (603) 6156 8708 Fax: (603) 6156 8707
Email: info@rangepharma.com
Website: www.rangepharma.com